

- [16] Wibowo EAP, Mauliani F, Nofitasari U, Setyani A, Widiarti N. 2017. Comparative analysis of chemical components of purified essential oil from nilam plants using Gas Chromatography. *Journal of Pure Applied Chemistry Resources*. 2017; 6(1):1-6. <https://doi.org/10.21776/ub.jpacr.2017.006.01.270>.
- [17] Kusuma HS, Mahfud M. The extraction of essential oils from patchouli leaves (*Pogostemon cablin* Benth) using a microwave air-hydrodistillation method as a new green technique. *The Royal Society of Chemistry Advances*. 2017; 7(3):1336-1347. <https://doi.org/10.1039/C6RA25894H>.
- [18] Werner. *Organic Complex Compound*. John Wiley & Sons, New York. 2018.
- [19] Adam F, Siti Hana AB, Yusoff MM, Tajuddin SN. Molecular dynamic simulation of the patchouli oil extraction process. *Journal of Chemical & Engineering Data*. 2014; 59(2):183-188. <https://doi.org/10.1021/je3013292>.
- [20] Keerthiraj M, Bhowmik A, Saha S, Dutta A, Chawla G, Kundu A. Optimisation of patchoulol in the lipid-soluble concentrates of *Pogostemon cablin* using response surface methodology (RSM) coupled with genetic algorithms (GA). *Industrial Crops and Products*. 2022; 182:114826. <https://doi.org/10.1016/j.indcrop.2022.114826>.
- [21] Muhammad S, H. P. S. AK, Abd Hamid S, Danish M, Marwan M, Yunardi Y, Abdullah CK, Faisal M, Yahya EB. Characterization of bioactive compounds from patchouli extracted via supercritical carbon dioxide (SC-CO₂) extraction. *Molecules*. 2022; 27(18):6025-6039. <https://doi.org/10.3390/molecules27186025>.